



RANA HOSPITAL

*Come with **PILES!** Go with **SMILES!***

Patient Information Leaflet

Going Home After
Abdominal Surgery

Going home after Abdominal Surgery

Everyone is different, and our bodies all react and recover in different ways. There are few hard and fast rules about what you should or should not do once you go home. The important thing is to strike a balance - try not to be in too much of a hurry, but do try steadily to increase the amount that you do. Listen to your body and do only as much as feels comfortable.

Abdominal pain

It is not unusual to suffer gripping pains (colic) during the first weeks following removal of a portion of your bowel. The pain usually lasts for a few minutes and goes away between the spasms.

If you have severe pain lasting more than a few hours or have a fever and feel generally unwell within a few weeks of your operation date, you should contact the ward on the telephone numbers provided.

If you have hiccups that persist for more than a day please contact the ward for advice.

Exercise and moving

Walking is encouraged from the day following your operation. You should aim to gradually increase your level of physical activity after you go home. Start with a short walk, for about 10 minutes, two or three times a day, and increase the distance as you feel able. By about six weeks you should be able to walk for half an hour or more if you could do so before your operation. Always try to walk as upright as you can to avoid backache. You may find climbing stairs tiring or uncomfortable at first. Try to get out of the house each day if the weather permits.

Try to sit in a chair which is high enough to rise from easily without putting strain on your abdominal muscles. A low sagging chair or sofa can mean that you sit hunched up and then strain your abdomen trying to get up. Try to sit with your lower back well-supported, maybe with a small cushion or a rolled up towel at waist level.

Household chores can involve a lot of bending and stretching and you may find this uncomfortable. Try to get some help with household chores, at least for the first one or two weeks after you go home. You should avoid standing for long periods and movements that involve stretching, pushing or pulling. You should avoid lifting heavy weights at first (including vacuum cleaners, heavy shopping bags and children) and not lift anything heavier than six to eight pounds (3-4 kilos, e.g. a full three pint kettle). If you need to hold children, try to let them climb onto your lap while you are already sitting rather than lifting them up.

Aim to gradually increase your physical activity. By about six weeks you should be able to walk for half an hour or more (if you could do so before your operation).

Once you are fully recovered you can normally undertake most activities. Heavy physical activities such as the gym should not usually be undertaken for several months.

Your bowels

Your bowel habit may change after removal of part of the bowel and you may find it becomes either more loose or more constipated than previously. Make sure you eat regular meals three or more times a day, drink adequate amounts, and take regular walks during the first few weeks after your operation.

If constipation lasts for more than three days then taking a laxative is advised. If you are passing loose stools more than ten times per day for more than a few days please call the ward and ask for advice. If you have a stoma, your stoma specialist nurse will explain about your stoma before you go home.

Clustering of bowel movements may be improved by adding a fibre supplement and a glass of water to the end of each meal. Muscle strengthening exercises combined with dietary changes may help with urgency and stool incontinence. Imodium (Loperamide) is an over-the-counter anti-diarrheal medication. It also increases sphincter pressure. Carry a survival pack (baby wipes, barrier ointment, Imodium), take advantage of aisle seating and be aware of restroom locations.

Your stoma

If you have a stoma and you have any problems with your stoma after you go home, please contact your stoma specialist nurse, you will be given contact details before you leave hospital.

Passing urine

Sometimes after bowel surgery you may experience a feeling that your bladder is not emptying fully. This usually improves with time. It is also worth keeping an eye on the colour of your urine. If you are well hydrated you should be passing straw colour urine. If you find that your urine is darker it may be a sign that you are dry and usually you should drink more. If you feel you are not emptying your bladder fully or if you have excessive stinging when passing urine, please ring the ward as you may have an infection.

Rest and sleep

You will probably be surprised at how tired you feel when you first go home. In hospital you felt OK, now you feel exhausted. This is normal. You probably didn't realise how little you actually do while in hospital, not even making a cup of tea. Don't worry - your strength and stamina will gradually return. It is important to mix activities with some rest as needed, but do get out of bed each day and get dressed if possible.

Try to plan a rest time each day, preferably on your bed. Let your family and friends know that this is important and that they should avoid disturbing you if possible.

You may find sleeping at night difficult at first. This may just be because your normal routine has been disturbed. Or you may have some discomfort or restricted movement. It is not uncommon to still have some pain when you first go home. A mild painkiller, such as paracetamol, before you go to bed may help.

Work and other activities

The length of time you need to take off work will depend on the operation you have had and the type of work that you do. You may be able to return to work within two to four weeks after your operation. People with jobs that involve a lot of heavy manual work may need more time off than those with less active jobs, but even sitting at a desk all day can be very tiring after an operation. You may wish to consider going back to work part-time initially if this is possible. If you do heavy manual work you should not normally go back until at least six to eight weeks after your operation. It can be useful to ask the doctor in clinic when you can go back to work, when you return to the hospital for your post-operative check-up.

There is no set time for resuming sexual relationships. Many people take a couple of months before they feel comfortable about this, but it is really up to you to know when you feel ready.

Driving

You should not drive until you are confident that you can drive safely. You should not start to drive again until your strength and speed of movement are up to coping with an emergency stop. You should also make sure that you are not drowsy from any painkillers and that your concentration is good. Most people do not start to drive for at least two weeks, and some will take longer, depending upon the operation you have had. If you find the seat belt uncomfortable over your scar a folded towel between you and the belt may help.

It might be sensible to check that you are insured to drive after surgery before you start driving.

Hygiene and wound care

It is quite safe to get your wound wet (unless you have been specifically advised not to do so). Either a bath or shower is fine, but do not have the water too hot at first. There is no evidence that adding salt to bath water helps with healing, and it may well dry your skin. Do not use antiseptics, bubble bath or anything else in the bath water, at least until the wound has fully healed. If you feel a little unsteady at first, use a non-slip bath mat and try to have someone else in the house in case you have difficulty getting out of the bath.

Your wound will go through several stages of healing. To start with you may feel tingling, itching or numbness. It may feel lumpy as new tissue forms and there may be some pulling as it heals. You should not go swimming until the wound is completely healed; often waiting for several months is advisable.

It is not unusual for wounds to be slightly red and uncomfortable during the first one or two weeks.

Please telephone the ward if your wounds:

- ❖ Become inflamed or swollen
- ❖ Become increasingly painful
- ❖ Start to discharge fluid

Getting back to “normal”

Having an operation can be a stressful experience, physically and emotionally. When you first go home you are likely to feel tired and unwell for a while. Things will get better. Some people report that it takes them 3-6 months to feel completely back to their normal selves, others recover much more quickly. It is common to feel a bit low in the first weeks and to become frustrated that you cannot do everything that you would like to do. Be patient!

Diet

A balanced, varied diet is recommended. Try eating three or more times a day. You may find that some foods upset you and cause loose bowel motions. If that is the case you should avoid those foods for the first few weeks following your surgery.

Some general advice is to:

- ❖ Sip fluids slowly.
- ❖ Avoid large amounts of caffeine as this can loosen stool output.
- ❖ Chew all foods thoroughly.
- ❖ Skipping meals may worsen watery stools and cause increased gas.
- ❖ Add new foods to your diet one at a time until their effect on bowel movements can be determined.

If you are finding it difficult to eat it is still important to obtain an adequate amount of protein and calories to help your body heal. You may benefit from having three to four nourishing, high protein, high calorie drinks such as Build-up or Complan (available in supermarkets and chemists) to supplement your food. If you are suffering from diarrhoea then it is important to replace the fluid loss and to drink extra fluid. Please call Rana Hospital for advice.

If you are losing weight without trying to or are struggling to eat enough, you may benefit from a consultation with the dietitian; ask your GP to refer you.

Summary

To enhance your recovery you will be expected to return to normal as quickly as possible. This means that you need to actively participate in your recovery by walking, eating and drinking. Each day you should feel some improvement but do contact the ward if you are worried about something.



RANA HOSPITAL

Come with PILES! Go with SMILES!

Contact details

Rana Hospital, Sirhind, Punjab (India)

Phone: 01763-220067

Email: ranapiles001@yahoo.in

www.ranapileshospital.com

Author: Rana Hospital

Review date: May 2014